

中文題目：主動脈-腸道瘻管引發之腸胃道出血：案例分享

英文題目：Suspect Aorto-Enteric fistula related Gastrointestinal Bleeding : A Case Report

作者：葉雙儀¹，張維安²

服務單位：高雄醫學大學附設中和紀念醫院¹內科部²胸腔內科


Case Presentation

This 75-year-old male has the history of gastric ulcer with Helicobacter pylori infection and gout. He suffered from hematochezia and hematemesis for several days. He was referred to our medical intensive care unit for suspecting gastrointestinal bleeding related hypovolemic shock.

Esophagogastroduodenoscopy revealed gastric ulcer without bleeder but much blood clot on the gastric fundus.

Computed tomography was done for palpable mass with pulsation over left lower abdomen. It showed huge

abdominal aortic aneurysm (AAA). His gastrointestinal bleeding much improved after medical treatment. However, emergent aortic stent implantation was done for recurrent hypovolemic shock episode. The hypovolemic shock also accompanied with massive lower gastrointestinal bleeding. After placing aortic stent, his hypotension and lower gastrointestinal bleeding much improved immediately.



Discussion

The clinical manifestation of ruptured abdominal aortic aneurysm is variable with respect to symptoms and time course. Gastrointestinal bleeding presented at abdominal aortic aneurysm rupture is uncommon case.